

**Daniel Atori**  
MINNA

**A**t least, three people were reportedly killed and 28 abducted by bandits in Farin-Shinge, Kulho, Jigawa, Dogo Fadama communities in Mashegu Local Government Area of Niger State.

The hoodlums, who attacked the communities Friday and Saturday, were said to have demanded seven a 50-litre jerry-can of petrol, Viju milk and other drinks for the release of the

**Lawrence Olaoye**  
ABUJA

President Muhammadu

## **Insecurity: Bandits kill 3, abduct 28 in Niger**

### **... demand petrol, Viju Milk, other drinks ...15 people abducted – Police**

abductees from the Farin-Shinge community

The Commissioner of Police, John Kuryas, said the bandits entered Mashegu village through Jigawa and rustled some cattle.

According to him, they broke a provision shop at Kawo village and also attacked Kulho village, a re-

mote community through Ibbi and abducted about 15 people.

Kuryas in a statement by the Police Public Relations Officer, DSP Wasiu Abiodun, said a manhunt had been launched for the criminals by police officers assisted by vigilantes to rescue the victims.

## **Buhari declares full-scale war against bandits, terrorists in Niger**

to express sincere sympathy for the government

bility of every member of the community and only

*New Telegraph* learnt that the miscreants attacked Farin-Shinge on the

outskirts of Kontagora at about 1.45pm when Jumaat (Friday) prayers were about to be held and abducted eight people.

## **Movement could slow early Parkinson's**

**Appolonia Adeyemi**

**P**eople with early-stage Parkinson's disease who regularly got one to two hours of moderate exercise twice a week, like walking or gardening, may have less trouble balancing, walking and doing daily activities later.

These are the results of a new study published in

the January 12, 2022, online issue of the journal 'Neurology.'

Parkinson's disease is a progressive nervous system disorder that affects movement. Symptoms start gradually, sometimes starting with a barely noticeable tremor in just one hand. Tremors are common, but the disorder also commonly causes stiffness or slowing of movement.

**50**